



Antioxidant drug, antifungal, anticancer, pharmacognosy, phytochemistry and pharmacological studies of *Cassia javanica* Linn. Plant

R B Singh

Scientist 'C' (UGC), Department of Zoology, School of Life Sciences, Dr. Bhimrao Ambedkar University, Khandari Campus, Agra, Uttar Pradesh, India

Abstract

Cassia javanica Linn. Plant is a small and medium sized tree, belongs to the plant family Leguminosae (Fabaceae). It blossoms in the spring season, mainly from the month of April to September. It is commonly known as the apple blossom cassia, pink lady, apple blossom shower and java shower. It is widely cultivated in the tropical regions of Asia. The plant consists of descending branches and it produces mass of gorgeous flowers. The fruits ripen in the fall. Studies have shown the presence of various phytochemical constituents mainly anthraquinone glycosides, flavonoids, alkaloids, sterols, tannins, saponins and reducing sugars in different parts of the plant. *Cassia javanica* Linn. Plant possesses various pharmacological activities antidiabetic, antioxidant, anticancer and antimycotic activities. It has long been used in traditional medicine to cure various diseases. It has been used as antipyretic, laxative and antimalarial drug. It is known to decrease virulence of pathogenic organisms. It is used in the treatment of gastric pain and cold. Due to complex nature of the disease there is a worldwide need for an ideal drug therefore, there are many antidiabetic herbs recommended in traditional medicaments. The presented review summarizes the information concerning the morphology, chemical constituents, uses and pharmacological activities of the plant.

Keywords: Antioxidant, antifungal, phytochemistry, anticancer, antidiabetic, anthraquinone glycosides, *Cassia javanica* Linn. Plant

Introduction

Leguminosae is the third largest family of angiosperms. It comprises of 740 genera which have a total of 19400 species. Ranging in habit from annual herbs to large trees, the leguminosae-family is well distributed throughout temperate and tropical regions of the world. *Cassia javanica* Linn. Plant is an important member of this plant Leguminosae- family. It has several common names such as pink cassia, pink shower, rainbow shower and white shower due to the presence of typical pink to crimson flowers. *Cassia javanica* Linn. Plant is distributed naturally from India to Malaysia, Sumatra, Indonesia, Southern China and the Phillipines. It is also cultivated in the tropical regions of Asia. It is known to be used as antidiabetic, anticancer, antifungal and antioxidant drug [1]. *Cassia javanica* Linn. Plant grows fast and reaches upto 25 meters height. Trunk diameter of the tree may near 35 cm.

The leaves are paripinnate, made up of 12 pairs of leaflets. Pale vose to crimson. Flowers are arranged in racemes. It produces typical hanging, thin, cylindrical legumes (pods) which are dark colored. The legumes are 30 to 40 cm long and upto 1.5 em in diameter. Each fruit contains numerous seeds (6.5 mm to 8.9 mm long and 5.6 mm to 7 mm wide). The seeds are obovate in shape and ventrally flattened. The seed coat is dark brown in color and has characteristic fracture lines [1].

The taxonomical status of the *Cassia javanica* Linn. Plant has been documented as shown in Table -1.

Vernacular Names

Hindi: Javaniki- Rani [2]

English: Java Cassia [3]

Tamil: Kondrai [4]

Table 1: Taxonomical Status of *Cassia javanica* Linn. Plant.

Kingdom	Plantae
Subkingdom	Tracheobionta
Superdivision	Spermatophyta
Division	Magnoliophyta
Class	Magnoliopsida
Subclass	Rosidae
Order	Fabales
Family	Fabaceae (Leguminosae)
Genus	<i>Cassia</i> L.
Species	<i>Cassia javanica</i> Linn.

Synonyms

Cassia Agnes (de Wit) Brenan

Cassia bartonii F.M. Bailey

Cassia nodosa Roxb.

Microscopy: The microscopic study of *Cassia javanica* Linn. Leaves revealed the presence of secondary vascular bundles with pericyclic fibres present in separate regions outside the phloem. The cortical sclerenchyma is absent. Cluster crystals are confined to cortical cells and dense multicellular trichomes are also present [5].

Phytochemistry: *Cassia javanica* Linn. Plant is a rich source of numerous phytoconstituents the particularly anthraquinones. The phytoconstituents are found in almost all parts of the plant. The constituents associated with individual plant organs have been mentioned in Table- 2.

Table 2: Phytoconstituents of *Cassia javanica* Linn. Plant

Plant Part	Secondary Metabolites
Leaves	Reducing sugars, proteins, alkaloids, tannins, glycosides, sterols, flavonoids, and saponins [3]; heptacosane, octacosane, hentriacontane, ceryl alcohol, octacosanol and hentriacontanol [6]. 1,3,4,6-tetrahydroxy-5-dimethoxy-2- methylanthraquinone and 1,3,5,8-tetrahydroxy-6-methoxy-2- methylanthraquinone; quercetin, emodin, chrysophanic acid, aloë-emodin, chrysophanol, physcion and its glucoside [7]. Nonacosane, triacontane, behenic acid, β -amyryn, emodin, kaempferol-3-O- β -D-glucosyl-6-O- α -L- rhamnopyranose; methyl mehenate, tetracosane, javanin [8]. Ent- Epiatzelechin-(4 α -8)-epiazzelechin [9]. Flavone rhamnoside 'javanin' [10].
Heartwood	Hentriacontane, β -sitosterol, cerotic acid and β -sitosterol-D-glucoside [10].
Stem bark	n-octacosanol and β -sistosterol [6]; 1,2-dihydro-1,3-dihydroxyl-6,8- dimethoxy-2-methylanthraquinone; 1,3,5,8-tetrahydroxy-6-methoxy-2- methylanthraquinone; 1,3,4,6-tetrahydroxy-5,8-dimethoxy-2- methylanthraquinone; 1,4-dihydroxy-6,7,8-trimethoxy-2- methylanthraquinone; 1-hydroxy-3,6,7,8-tetramethoxy-2- methylanthraquinone; 4,4'bis(1,5-dihydroxy-7-hydroxymethyl-2-methyl-3-Methoxy) anthaquinone [7].
Seeds	Chrysophanol, physcion; 1,5-dihydroxy-4,7-dimethoxy-2- methylanthraquinone-3-O- α -rhamnopyranoside; 1,3,6,7,8-Pentahydroxy-4-methoxy-2-methylanthraquinone and galactomannans [11].
Flowers	Flavonoid glycosides (leucocyanidin-4'-O-methylether-3-O- β -d- galactopyranoside; dihydrorhamnetin-3-O- β -d-glucopyranoside; quercetin-3', 4', 7-trimethylether-3-O- α -l-rhamnopyranoside; kaempferol-3- rhamnoglucoside) alongwith quercetin [12]. 5, 3-dihydroxy-2-phenyl-4- chrome-7, 4-digulucoside; naringenin-7-glucoside; 3-glucoside of pelargonidin. Paeonidin-3-O- α -L(-)-hannopyranoside [8] and Glactomannans [13].
Root	Emodin-8-rhamnoside, 5-hydroxyemodin-8-rhamnoside; 1, 3- dihydroxy-5, 6, 7-trimethoxy-2-methylanthraquinone; 1, 4-dihydroxy-8-methoxy-2- methylanthraquinone-3-O- β -D-glucopyranoside; 1, 8-dihydroxy-6, 7- dihydroxy-2-methylanthraquinone [7]. 3',6' dihydroxy-4- methoxychalcone, 1,6-dihydroxy~ and 1,5,6-trihydroxy-3- Methylanthraquinone. 8-O- α -L-rhamnopyranoside. [8].
Bark	Proanthocyanidins containing flavan-3-ol (epiazzelechin and epicatechin) [14].

Structure of Anthraquinone

The anthraquinones are found in *Cassia javanica* Linn. Leaves in free as well as glycosylated form. Additionally, Leaves contain sugars, flavonoids and isoprenoid

compounds. Roots, heartwood and seeds are rich in anthraquinones. Flowers contain a variety of flavonoids in free and glycosylated forms. And structure are shown in Figure-1.




Fig 1: Structure of Anthraquinone from *Cassia javanica* Linn. Plants parts

Pharmacological studies

Cassia javanica Linn. Plant possesses variety of pharmacological actions some of which are as follows:

Hypoglycaemic activity: The aqueous ethanolic and chloroform extracts of leaves showed hypoglycaemic activity on laboratory-bred male Wistar albino adult rats [3].

Anticancer and antimycotic activity: The anticancer and antimycotic activities of honey samples from flowers of Egyptian *Cassia javanica* Linn. Plant against colon, breast and liver tumor cell lines (HCT-116, HTB-26 and HepG2) and over clinical dermatophytes (*Tricophyton*, *Microsporum* and *Epidermophyton*) have been evaluated [15].

Antioxidant activity: Antioxidant potential of methanolic extract of leaves was evaluated in superoxide anion radical scavenging assay and it revealed a strong antioxidant potential of 50.4% at a concentration of 300 μ g/ml [16].

Antimicrobial activity: Propelargonidin from *Cassia javanica* Linn. Plant is fungitoxic [17]. *Cassia javanica* Linn. Is fungitoxic to *Rhizoctina solani* [18].

Antiviral activity: Ent-epiazzelechin-(4 α +8)-epiazzelechin (EEE) found in the fresh leaves of *Cassia javanica* Linn. was subjected to *in vitro* testing against anti-Herpes Simplex Virus-2 (HSV-2) using 2,3-Bis-(2-methoxy-4-nitro-5-

sulfophenyl)-2H-tetrazolium-5-carboxanilide (XTT) and plaque reduction assays. EEE was found to prevent penetration of cells by HSV-2 and also negatively affected HSV-2 replication in a dose dependent manner [9]. The saline extract of *Cassia javanica* Linn. Seeds showed

haemagglutinating and haemolytic activity against RBCs [10]. And structure of haemolytic from seeds are given in Figure-2.

Haemolytic activity




Fig 2: Structure of Haemolytic activity of *Cassia javanica* Linn. Seeds

Conclusion

Cassia javanica Linn. Plant is an important medicinal plant of Family Leguminosae. It belongs to *Cassia*, a genus known for anthraquinones with laxative/purgative action. The plant or its extracts have exhibited numerous activities (such as hypoglycaemic, anticancer, antiviral and antimycotic actions) possibly due to the presence of flavonoids, anthraquinones and sterols. Elementary. Studies suggest a therapeutic potential exists for this plant beyond the traditional ages.

References

- Rocas NA. *Cassia javanica* Linn. Part II: Species descriptions, 2012, 372–374.
- Pant MH, Sharma N. *Medicinal Plants Research*, 2010;4(20):2144–2147.
- Kumavat UC, Shimpi SN, Jagdale SP. *Journal of Advanced Pharmaceutical Technology & Research*, 2012;3(1):47–51.
- Ganesan S. *Natural Product Radiance*, 2008;7(2):166–172.
- Shaheen ASM. *Turkish Journal of Botany*, 2007;31:297–310.
- Joshi KC, Tholia MK, Sharma T. *Planta Medica*, 1975;28(2):190–192.
- Dave H, Ledwani L. *Indian Natural Products and Resources*, 2012;3(3):291–319.
- Ganapaty S, Thomas PS, Ramana KV, Vidyadhar K, Chakradhar V. *Natural Remedies*, 2002;2(2):102–120.
- Cheng HY, Yang CM, Lin TC, Shieh DE, Lin CC. *Medical Microbiology*, 2006;55(2):201–206.
- WOI. *The Wealth of India: A Dictionary of Indian Raw Materials and Industrial Products*. Revised ed, 1992, 345–346.
- Tiwari RD, Sharma MN. *Planta Medica*, 1981;43(4):381–383.
- Tiwari RD, Yadava OP. *Phytochemistry*, 2001;10(9):2256–2263.
- Andrade CT, Azero EG, Luciano L, Gonçalves MP. *International Journal of Biological Macromolecules*, 1999;26(2–3):181–183.

14. Kashiwada Y, Iizuka H, Yoshioka K, Chen RF, Nonaka G, Nishioka I. *Chemical and Pharmaceutical Bulletin*,1990;38(4):888-893.
15. El-Gendy MMA. *Journal of Microbial Biochemistry and Technology*,2010;2:118-123.
16. Kaur P, Arora S. *Chinese Clinical Medicine*,2010;5(8):57.
17. WOI. *The Wealth of India: A Dictionary of Indian Raw Materials and Industrial Products. A-Ci.* 2004;1:224-225.
18. Punyasiri PAN, Tanner GJ, Abeyasinghe ISB, Kumar V. *Phytochemistry*,2004;65:2987-2994.